



INSTRUCTIONS FOR COLONOSCOPY Dr Anderson Patients

Colonoscopy: The visual examination of the large intestine (colon) using a lighted flexible fiber optic endoscope.

Note: Individual responses to laxatives do vary. This prep may cause multiple bowel movements. Often works within 30 minutes but may take as long as 3 hours. If you are unable to keep the prep down, please call your physician.

Consult your Primary Care Physician if you are taking or have the following medical concerns:

Coumadin/Jantoven/warfarin: Stop it five (5) days before your colonoscopy if approved by your physician prescribing this medication. **Do not take this medication on the day of your procedure. Ask your physician when to resume this medication**
Plavix (clopidogrel)/Pradaxa/Pletal (cilostazol)/Effient and Eliquis; Stop it three (3) days before colonoscopy-**only if approved by prescribing physician**

Brilinta: Stop five (5) days before colonoscopy- **only if approved by prescribing physician**

Aggrenox: Stop three (3) days to five (5) days prior to colonoscopy- **only if approved by prescribing physician**

Xarelto/Savaysa: Stop 48 hours before colonoscopy- **only if approved by prescribing physician**

Iron pills: Stop for seven (7) days before colonoscopy

Dialysis Treatment: Alternate treatment may be used.

****Anti-Seizure Medication:** Take at least 2 hours before the scheduled colonoscopy with a small sip of water.

****Blood Pressure/Heart Medication:** take your morning dose at least 2 hours before your colonoscopy with a small sip of water.

Insulin: You may need to adjust your insulin dosages. Please speak with your monitoring physician for further instructions. **Ask your physician when to resume your insulin.**

Inhalers: Please bring your "rescue" inhaler to your appointment, even if you have not needed to use it for some time.

Heart Valve Replacement/ Pacemaker with Internal Defibrillator

No tomatoes or corn 2 days prior to procedure

APPROVED CLEAR LIQUIDS LIST

Water

Jell-O – any flavor – no red Jell-O

Coffee or Tea without creamer

Strained fruit juices without pulp

Clear sodas – 7up, Sprite, Ginger Ale, etc.

Ensure and Ensure plus (vanilla only)

Broth or Bouillon

Popsicles – no red or purple

No red or purple liquids

DAY OF COLONOSCOPY

1. Nothing by mouth after midnight except approved **medications listed above**** with a sip of water. You may brush your teeth and rinse with water. **NO GUM, MINTS OR LOZENGES. NO CHEWING TOBACCO**
2. **You will be medicated and you need to bring someone with you to drive you home.**
3. Bring a current list of medications and colostomy supplies if applicable; if Menstruating please use a tampon if possible.

On the day of your procedure, report to the following unless otherwise instructed by your physician:

The Endoscopy Center of Topeka, 785.354.1254

2200 SW 6th Avenue, Suite 103

Topeka, KS 66606

Check-in Date: _____

Check-in Time: _____

KMC Gastroenterology, 785-354-8518

If you are unable to keep your appointment, please notify the office within 48 hours prior to your scheduled procedure